

Charge Ordering in the $Pt_2(dta)_4I$ Compound

Marie-Bernadette LEPETIT^{1,*} and Vincent ROBERT¹

¹*Laboratoire de Physique Quantique, IRSAMC/UMR 5626, Université Paul Sabatier,
118 route de Narbonne, F-31062 Toulouse, Cedex 4, FRANCE*

(Dated: February 2, 2008)

The present paper tries to elucidate the electronic structure of the $Pt_2(L)_4I$ family of compounds in the semi-conducting and metallic phases, using correlated *ab initio* embedded fragment calculations. The usual model associating the different phases with different intra-dimer charge orders, is energetically ruled out by our results. A careful analysis of the local low-energy degrees of freedom lead us to propose a mechanism associating charge transfer from the iodine $5p_z$ orbital with the vibrational degrees of freedom of the *dta* ligands. It is shown that such a model can account for both the XPS and IR experimental results at the phase transition.

Low-dimensional compounds have attracted a lot of interest in both experimental and theoretical communities for the last twenty years. Among these, the one-dimensional (1D) halogen-bridged transition-metal complexes exhibit several very peculiar properties which have induced a lot of research work. One can cite, for instance, large third-order non-linear optical properties [1], photo-generation and annihilation processes of mid-gap states [2], or intense dichroic inter-valence charge transfer absorption [3]. These systems are perfectly one-dimensional, mixed valence and strongly correlated, being thus paradigms of strongly interacting 1D systems, and exhibiting a large number of different phases. The nature of the ground state can be controlled not only by pressure and temperature but also by chemical substitution of either transition metal atoms, halogen atoms or even counter ions [4]. However all the so-called MX (M metal, X halogen) systems which have been synthesized so far are insulators or semi-conductors. Extensive works have been done without success, on these systems, in order to induce metalization under pressure [5]. Therefore researches have switched focus toward the synthesis of bi-nuclear [6] (and very recently, tetra-nuclear [7]) MMX systems. The $Pt_2(dta)_4I$ complex (*dta*=dithioacetate= $CH_3CS_2^-$) has thus become the second example (following $KCP(Br)$ [8]) of 1D system exhibiting metallic transport without involving a π electronic system. This system has since then become the archetype of a series of similar MMX intriguing compounds.

The $Pt_2(dta)_4I$ compound has a neutral chain structure consisting of dimeric units $[Pt_2(dta)_4]$ bridged through halides (see figure 1). From electrical conductivity and thermoelectric power measurements [9] three different phases have been identified in the range of temperatures from 90K to 450K. The system is a semi-conductor between 90K and $T_{MS} = 300K$. From 300K to 373K, the system is metallic and then undergoes a metal to metal phase transition at $T_{MM} = 373.4K$.

Let us now analyze the electronic structure. A formal charge analysis leads to the $(Pt)_2^{5+}(dta)_4^{4-}I^-$ charge transfer form. Ligand field analysis shows that the Fermi




FIG. 1: Crystal structure of the $Pt_2(dta)_4I$ compound at 298K [6]. One should note that the two PtS_4 units of the $Pt_2(dta)_4$ dimer are twisted from the eclipsed position with an average angle of 21° .

level orbitals are the Pt non-degenerated $5d_{z^2}$ orbitals, z being the axis of the chains. It results that the low-energy physics of the $Pt_2(dta)_4I$ compound can be described as a correlated three-quarter filled band supported by the platinum $5d_{z^2}$ orbitals. Such a mixed-valence system can be subject to a large number of valence instabilities associated with different charge orderings, namely one can find

- (A) $I^- \dots Pt^{2.5+} \dots Pt^{2.5+} \dots I^- \dots Pt^{2.5+} \dots Pt^{2.5+} \dots I^-$
- (B) $I^- \dots Pt^{2+} \dots Pt^{3+} \dots I^- \dots Pt^{2+} \dots Pt^{3+} \dots I^-$
- (C) $I^- \dots Pt^{2+} \dots Pt^{3+} \dots I^- \dots Pt^{3+} \dots Pt^{2+} \dots I^-$
- (D) $I^- \dots Pt^{3+} \dots Pt^{3+} \dots I^- \dots Pt^{2+} \dots Pt^{2+} \dots I^-$

Based on several experimental results, different authors have tried to interpret the electronic structure of the different observed phases according to the above charge orderings.

In the 150 – 300K temperature range, XPS experiments [9] exhibit two signals associated respectively with Pt^{2+} and Pt^{3+} . Similarly, IR [9] spectra show a doubling of the $C = S$ stretching mode upon cooling from the metal (M) to the semi-conductor (SC) phase. This double peak has been interpreted as a signature of a double valency of the platinum atoms in the SC phase, therefore excluding the electronic structure pictured in (A). As for magnetic susceptibility measurements, a slight increase is observed upon cooling, below T_{MS} . Kitagawa *et al* [9]

therefore concluded to a zero spin gap and a $4k_F$ -CDW Mott-Hubbard ground state for the SC phase, that is an electronic structure of type (B). However, it should be pointed out that i) the XPS experiments have a very short time scale ($\sim 10^{-17}$ s) and therefore cannot differentiate between mixed and averaged valencies (as stated in references [9] and [?]), ii) the XPS Pt^{2+} and Pt^{3+} signals intensities exhibit a 3 to 1 ratio, incompatible with any of the above-cited charge orderings.

The aim of this paper is to analyze the above issue, using *ab initio* quantum chemical calculations. We used for this purpose the embedded fragment technique associated with *ab initio* spectroscopy methods, that is large atomic basis sets [12] and configurations interaction techniques dedicated to the accurate evaluation of both transitions energies and states wave-functions. More precisely we used the Difference Dedicated Configuration Interaction (DDCI) method [13, 14]. It this method the orbital space is split into three sets : **the occupied orbitals** that are always doubly-occupied in the system, **the active or magnetic orbitals** that have fluctuating occupation numbers or spins in the ground and excited states, **the virtual orbitals** that are always empty in the system. The DDCI method is an exact diagonalization method on a selected configuration space designed as follow :

- all configurations that can be built from the above occupation rules, this is the Complete Active Space (CAS) which is used as a reference space and corresponds to the support of the model Hamiltonian,
- then on this CAS, all single excitations and all double excitations participating in the excitation energy at the second order of perturbation (that is the configurations responsible for the screening effects on the interactions within the CAS).

This approach has been very successful in the quantitative evaluation of quantities such as magnetic couplings, charge order, spin polarization, etc. both in molecular magnetic systems and in strongly correlated materials (see for instance references [15, 16, 17]). Its accuracy can be exemplified on the high- T_c superconductor Nd_2CuO_4 for which the effective exchange coupling has been computed within experimental accuracy ($J_{calc} = -126.4$ meV [16], $J_{exp} = -126 \pm 5$ meV [18]). The success of these embedded fragment methods in transition metal compounds is due i) to the locality of the d Fermi electrons responsible for the low-energy properties [19] and ii) to the possibility of using quantum chemical methods able to accurately treat open-shell problems and screening effects.

The fragment used in the present calculations is composed of one $Pt_2(dta)_4$ dimeric unit and its neighboring iodine atoms, that is $IPt_2(dta)_4I$. Its geometry has been extracted from the X-ray structure at room temperature [6]. This fragment was embedded in a large bath of

total-ion pseudo-potentials [12] (TIP) and point charges, designed in order to account for the Madelung potential and the short-range exclusion and exchange effects of the rest of the crystal. We have considered the three different possible fillings for the platinum dimer, i.e. $(Pt_2)^{6+}$, $(Pt_2)^{5+}$ and $(Pt_2)^{4+}$ involved in the charge orderings (A) to (D) — that is $(IPt_2(dta)_4I)^{0,1-2-}$.

Results on the $(Pt_2)^{n+}$ systems ($n = 4, 5, 6$) exhibit a very low-lying bonding $\sigma_d = (d_{z^2}(Pt_1) + d_{z^2}(Pt_2)) / N_\sigma$ orbital (where 1 and 2 refer to the two Pt atoms of the dimer and N_σ is the normalization constant) and a Fermi level anti-bonding $\sigma_d^* = (d_{z^2}(Pt_1) - d_{z^2}(Pt_2)) / N_{\sigma^*}$ orbital. These two orbitals are separated by the (dta) -ligand bonding and non-bonding orbitals, as well as all the other $Pt(5d)$ orbitals combinations.

$(Pt_2)^{5+}$. The intra-dimer charge localization within the $(Pt_2)^{5+}$ dimer, as pictured in modes (B) and (C), necessitates the excitation of an electron from the σ_d orbital toward the σ_d^* one. Indeed, $\sqrt{2} |d_{z^2}(Pt_2)d_{z^2}(Pt_1)\overline{d_{z^2}(Pt_1)}} = |\sigma_d\sigma_d^*\overline{\sigma_d}| + |\sigma_d\sigma_d^*\overline{\sigma_d^*}|$. However, the $\sigma_d \rightarrow \sigma_d^*$ excitation requires quite large an energy ($\simeq 5$ eV) and is therefore strongly hindered. Coherently, we found the electronic density to be highly delocalized between the two platinum atoms on the $(Pt_2)^{5+}$ system, despite the twisting of the (dta) ligands which is usually considered as responsible for intra-dimer charge localization $Pt^{2+}Pt^{3+}$ [20]. Indeed, the differential Mlliken population between the two Pt atoms is $ch(Pt_1) - ch(Pt_2) = 0.02\bar{e}$ (in full agreement with our periodic mean-field calculations based on the experimental crystal structure [6] which yield $0.03\bar{e}$). These results therefore exclude the possibility of a charge localization within the platinum dimer. In conclusion, the description of the electronic structure of the semi-conducting phase as intra-dimer charge ordered (whether following the (B) or (C) patterns) is incompatible with *ab initio* calculations.

From the above results we see that the bonding σ_d orbitals on the $(Pt)_2(dta)_4$ dimers are not pertinent for the low-energy physics of the compound. The pertinent model should be a half-filled one-band (extended) Hubbard model (EHM) based on the dimer σ_d^* orbitals. Such a model would be in agreement with the thermoelectric power experiments. The physics of the one-dimensional EHM is well known [21] and unfortunately does not exhibit any metallic phase for positive one-site repulsion U and first neighbor repulsion V . One should therefore invoke other degrees of freedom than the one supported by the d_{z^2} orbitals of the platinum atom in order to understand the different phases of this compound.

Let us first examine the low-energy electronic degrees of freedom of the $(Pt)_2(dta)_4$ dimer. The first excited state of both the $(Pt_2)^{5+}$ and $(Pt_2)^{6+}$ dimers is a ligand to metal single excitation where a non-bonding π electron of the ligand is excited into the σ_d^* orbital. The computed excitation energies are 2.01eV and 0.76eV for the 5+ and

6+ systems respectively.

The low-energy physics will be governed by the energy scale of the inter-dimer interactions. Even-though the $(Pt)_2(dta)_4 I (Pt)_2(dta)_4$ system is much too large to be investigated using the DDCI methods, the energy scale of the inter-dimer interactions can be evaluated using qualitative methods. Periodic Hartree-Fock calculations yield a band width of $0.79 eV$. The coulombic repulsion of two electrons on the dimer σ_d^* orbital $U_{\sigma_d^*}$ can be roughly evaluated by computing

$$E((Pt_2)^{4+}) + E((Pt_2)^{6+}) - 2 E((Pt_2)^{5+})$$

which yields $U_{\sigma_d^*} \simeq 3.9 eV$. It results that the effective hopping t between the σ_d^* orbitals of two dimers should be of the order of magnitude of $0.79/4 \simeq 0.2 eV$, while the effective exchange should be of the order of $4t^2/U_{\sigma_d^*} \simeq 0.04 eV$. These estimations completely scale out the possible mobilization of the $2.01 eV$ first state on $(Pt_2)^{5+}$. The $0.76 eV$ first state on $(Pt_2)^{6+}$ is also excluded since it is of a different approximate symmetry than the local ground state (π instead of σ) and thus cannot couple in a significant way with the low-energy band.

The only other possible electronic degree of freedom involves the transfer of a $5p_z$ electron of the iodine toward the σ_d^* orbital of the dimeric unit. The possible importance of such excitations has already been pointed out by S. Yamamoto [22]. In order to accurately check their pertinence, one should be able to evaluate the iodine $5p_z$ to σ_d^* excitation energy and to compare it to the effective transfer integral between these orbitals. Unfortunately it is impossible to accurately evaluate these values using *ab initio* methods, since one should be able to compute excitations on the $I(Pt)_2(dta)_4 I (Pt)_2(dta)_4 I$ system, a presently out of reach problem. However, it is possible to get a rough estimate of the $\varepsilon_{\sigma_d^*} - \varepsilon_I$ orbital energy difference between the σ_d^* orbital and the p_z orbital of the iodine, as well as of the transfer integral between the iodine $5p_z$ orbital and the dimer σ_d^* orbital using mean-field Hartree-Fock theory. It comes $\varepsilon_{\sigma_d^*} - \varepsilon_{p_z} \simeq 0.79 eV$ and $t_{\sigma_d^*, p_z} \simeq 0.56 eV$. From these parameters it is possible to compute, using a simple two-bands tight-binding model, the charge transfer from the iodine $5p_z$ orbital toward the σ_d^* orbital of the dimer. The computed charge transfer is quite large with 0.89 electron. The correlation effects, that are expected to be quite important in these systems, will probably reduce this value. However, it is reasonable to still expect a significant charge transfer between the two orbitals.

One of the important consequences of such a two-band model is that it becomes easy to explain the XPS experimental results for the $Pt_2 I(dta)_4$ compound. Indeed, the total charge carried by the platinum atoms is now allowed to fluctuate. An average electron transfer of η , from the iodine to the platinum dimer will change the

average charge of the latter from 2.5 to $2.5 - \eta$, in consequence of which the XPS experiments will see $0.5 + \eta Pt^{2+}$ and $0.5 - \eta Pt^{3+}$ ions. The experimental ratio between the Pt^{2+} and Pt^{3+} signals, which is close to 3 : 1, may be understood along this picture with $\eta \simeq 1/4$. Such a value corresponds to an electronic structure mixing in equal weights, i) the configuration where the iodine is always doubly occupied and ii) the set of singly-excited determinants where an electron has been transferred from an iodine atom toward a platinum dimer.

We have examined all electronic low energy degrees of freedom, let us now examine the effects of the vibrational degrees of freedom. One of us (VR) suggested, from extended Hückel calculations, that the internal vibrational degrees of freedom of the $Pt_2(dta)_4$ dimers — and in particular the (dta) torsion modes — must be of importance in the electronic structure of the system [20]. These vibrational modes are essentially local and independent. Thus, they can be associated with optical phonons and modeled using Holstein coupling to the electronic Hamiltonian. The system can thus be reasonably represented by a two-bands, 3/4-filled, extended Hubbard-Holstein model.

The dta torsion modes expected to be important for the $Pt_2 I(dta)_4$ compound are soft modes with low vibrational frequencies. The phase diagram of the 3/4-filled, extended-Hubbard Holstein model have been extensively studied [23]. For low frequency vibrational modes, and weak to moderate electron-phonon coupling constant, it exhibits first a Luttinger liquid phase with renormalized parameters, then, for larger coupling, a metallic phase characterized by dominant $4k_F$ -Charge Density Wave (CDW) fluctuations and finally, for intermediate coupling, an insulating $4k_F$ -CDW phase, with an exponentially opening gap. The transition between the different phases is controlled by the ratio between the hopping integral renormalized by the Franck-Condon factors, and the electronic repulsion renormalized by the phonons.

Starting from the $4k_F$ -CDW phase, not too far from the phase transition, and increasing the temperature, one excites the system in vibrational modes of higher quantum number and thus the Franck-Condon factors increase. Consequently, the effective hopping integral increases and the system enters the metallic phase, in agreement with the $Pt_2 I(dta)_4$ transport measurements.

In this model, the SC phase is characterized as a $4k_F$ -CDW, that is a phase presenting a charge alternation one unit cell out of two. The Pt_2 dimers can thus be expected to see a charge alternation $(Pt_2)^{2.5-\eta+\mu}$, $(Pt_2)^{2.5-\eta-\mu}$. Such a dimer charge alternation is in agreement with the doubling of the $C = S$ vibrational signal seen in IR experiments at T_{MS} .

In conclusion, accurate local *ab initio* calculations on the $Pt_2(dta)_4$ dimer have called in question the possibility of intra-dimer charge localization in the *MMX* compounds. Indeed, in the $Pt_2 I(dta)_4$ compound, the two

Pt atoms of a dimer are so strongly covalently bonded — due to short *Pt* – *Pt* distances — that intra-dimer charge localization can be considered as impossible. The only electronic degree of freedom of the dimers, pertinent for the low-energy physics, is the anti-bonding σ_d^* orbital which is delocalized on the two *Pt* atoms. In consequence the electronic structure of the different phases of the $Pt_2I(dta)_4$ compound cannot be understood in terms of different patterns of charge localization on the platinum atoms ((A) to (D)).

After a careful analysis of the different low-energy degrees of freedom, we have proposed a new model, that is able to account for both the *ab initio* results and the experimental ones. This model involves, in addition to the σ_d^* dimer orbital, i) the charge transfer excitation from the $5p_z$ orbital of the iodine and ii) the vibrational degrees of freedom of the *dta* intra-dimer ligands. Such a 3/4-filled, two-bands, extended-Hubbard Holstein model can account for the phase transition, as well as the XPS and IR experimental results.

Looking now at similar systems, the present conclusions should hold for analog $Pt(RCS_2)_4I$ (*R* = ethyl, propyl, butyl) compounds, since, both the intra-dimer distances are very similar (ethyl : 2.684Å, propyl : 2.675Å to 2.689Å, butyl : 2.68Å), and the external part of the intra-dimer ligand does not seem to play an important role on the electronic parameters [25].

For the $A_4Pt_2(pop)_4X$ family, the *Pt* – *Pt* distances are somewhat larger (~ 2.8 Å) resulting in a smaller bonding-antibonding intra-dimer splitting. As a consequence, the resistance of the intra-dimer delocalization to localizing perturbations should be weaker. However, experimental data do not exhibit such a localization, except maybe under large pressures where inter-dimer effects start to compete [24]. The pertinence of the charge-transfer degree of freedom from the iodine to the dimer is also questionable in these systems. Indeed, it is strongly dependent on the *Pt* – *Pt* and *Pt* – *I* distances. A sign that supports the hypothesis of the non-relevance of this degree of freedom for the *pop* systems is the 1 : 1, Pt^{2+}/Pt^{3+} ratio observed in the XPS spectra of the $A_4(Pt)_2(pop)_4X$ compounds.

We would like to thank D. Maynau for providing us with the CASDI codes and N. Guihry for helpful discussions.

* Electronic address: marie@irsamc.ups-tlse.fr

- [1] Y. Iwasa, E. Funatsu, T. Hasegawa and T. Koda, Appl. Phys. Letters **59**, 2219 (1991).
- [2] Y. Wada, N. Matsushita and M. Yamashita, Synth. Met. **56**, 3395 (1993).
- [3] H. Okamoto, T. Mitani, K. Toriumi and M. Yamashita, Phys. Rev. **B 42**, 10381 (1990).
- [4] N. Matsushita, Synth. Met. **56**, 3401 (1993).
- [5] See for instance : L.V. Interrabte, K.W. Browall and F.P. Bundy, Inorg. Chem. **13**, 1158 (1974) ; I. Shirotni *et al.*, Synth. Metal **64**, 256 (1994).
- [6] H. Kitagawa, S. Nakagami and T. Mitani, Synth. Metals **116**, 401 (2001).
- [7] K. Sakai *et al.*, J. Am. Chem. Soc. **120**, 8366 (1998).
- [8] J.-F. Thomas, Solid-State-Communications **42**, 567 (1982).
- [9] H. Kitagawa *et al.*, J. Am. Chem. Soc. **121**, 10068 (1999).
- [10] Y. Miyazaki *et al.*, J. Phys. Chem. **B 106**, 197 (2002).
- [11] G.A. Sawatzky and E. Antonides, Journal de Physique **C 4**, 117 (1976).
- [12] The inner-core electrons of the platinum ($[1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^{10} 4p^6 4d^{10} 4f^{14}] 5s^2$), iodine ($[1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^{10} 4p^6 4d^{10} 4f^{14}]$), sulfur ($[1s^2 2s^2 2p^6]$) and carbon ($[1s^2]$) atoms of the fragment are represented by effective core potentials (ECP). The outer electrons have been described in a basis set of triple-zeta plus polarization quality. ECP and basis set references are as follow : **Pt** : M. Casarrubios and L. Seijo, J. Chem. Phys. **110**, 784 (1999). **I** : A. Bergner, M. Dolg, W. Kuechle, H. Stoll, H. Preuss, Mol. Phys. **80**, 1431 (1993). **S** and **C** : A. Bergner, M. Dolg, W. Kuechle, H. Stoll and H. Preuss, Mol. Phys. **80**, 1431 (1993). **H** : W.J. Hehre, R.F. Stewart and J.A. Pople, J. Chem. Phys. **51**, 2657 (1969). The **TIP** used for the embedding are : **Pt** : S. Zurita, J. Rubio, F. Illas and J.-C. Barthelat, J. Chem. Phys. **104**, 8500 (1996), **I** : V. Robert, F. Spiegelman and M.-B. Lepetit, Chem. Phys. **287**, 217 (2003).
- [13] J. Miralles, J. P. Daudey and R. Caballol, Chem. Phys. Lett. **198**, 555 (1992) ; V. M. García, O. Castell, R. Caballol and J. P. Malrieu, Chem. Phys. Lett. **238**, 222 (1995); V. M. García, M. Reguero and R. Caballol, Theor. Chem. Acc. **98**, 50 (1997).
- [14] The present calculations have been performed using the MOLCAS and CASDI suite of programs. Molcas Version 5. K. Anderson *et al.*, Lund University, Sweden (2000).
- [15] J. Cabrero *et al.*, J. Phys. Chem. **A 104**, 9983 (2000).
- [16] See for instance : D. Munoz, F. Illas and I. de P.R. Moreira, Phys. Rev. Letters **84**, 1579 (2000), and references therein.
- [17] N. Suaud and M.-B. Lepetit, Phys. Rev. Letters **88**, 056405 (2002).
- [18] T.R. Thurston *et al.*, Phys. Rev. Letters **65**, 263 (1990).
- [19] I. de P.R. Moreira *et al.*, Phys. Rev. **B 59**, R6593 (1999) ; M.-B. Lepetit, Recent Research Developments in Quantum Chemistry 3, Transworld Research Network (2002).
- [20] S.A. Borshch, K. Prassides, V. Robert and A.O. Solonenko, J. Chem. Phys. **109**, 4562 (1998).
- [21] See for instance : J. Voit, Phys. Rev. **B 45**, 4027 (1992) ; M. Nakamura, Phys. Rev. **B 61**, 16377 (2000) ; M. Tsuchiizu and A. Furusaki, Phys. Rev. Letters **88**, 056402-1 (2002).
- [22] S. Yamamoto, Phys. Rev. **B 63**, 125124 (2001).
- [23] Ph. Maurel and M.-B. Lepetit, Phys. Rev. **B 62**, 10744 (2000) ; Ph. Maurel, M.-B. Lepetit and D. Poilblanc, Eur. Phys. J. **B 21**, 481 (2001) ; Ph. Maurel and M.-B. Lepetit, Phys. Rev. **B 66**, 075117 (2002).
- [24] S. Yamamoto, Phys. Rev. **B 64**, 140102 (2001).
- [25] T. Onishi, V. Robert and M.-B. Lepetit, to be published.